arXiv:1211.0219v2 [astro-ph.CO] 8 Nov 2012

USTC-ICTS-12-13

The quintessence field as a perfect cosmic fluid of constant pressure

Wenzhong Liu[*], Jun Ouyang[†] and Huanxiong Yang[‡]

Interdisciplinary Center for Theoretical Study,
University of Science and Technology of China,
Hefei, 230026, P. R. China

November 9, 2012

Abstract

We study the cosmology of a quintessence scalar field which is equivalent to a non-barotropic perfect fluid of constant pressure. The coincidence problem is alleviated by such a quintessence equation-of-state that interpolates between plateau of zero at large redshifts and plateau of minus one as the redshift approaches to zero. The quintessence field is neither a unified dark matter nor a mixture of cosmological constant and cold dark matter, because the quintessence density contrasts decay monotonously on sub-horizon scales and the squared sound speeds of quintessence perturbations do not vanish. What a role does the quintessence play is dynamic dark energy. Though the clustering of quintessence decays drastically, it could remarkably impact the growth rate of the density perturbations of non-relativistic matters at early stage.

---

[*] wzhliu@mail.ustc.edu.cn
[†] yangjun@mail.ustc.edu.cn
[‡] hyang@ustc.edu.cn

1 Introduction

Recent cosmic observations, including Type Ia Supernovae [1, 2], Large Scale Structure (LSS) [3, 4, 5, 6, 7, 8] and Cosmic Microwave Background (CMB) [9, 10, 11, 12] have independently indicated that the evolution of the universe is currently dominated by a homogeneously distributed cosmic fluid with negative pressure, the so-called dark energy. The dark energy fills the universe making up of order 74% of its energy budget. The remaining energy fraction in the present universe is about 22% occupied by the pressureless cold dark matter and 4% occupied by baryons. Despite many years of research and much progress, the nature and the origin of dark energy does still remain as an open issue.

Phenomenologically, the best candidate of dark energy that fits perfectly the observation data is the so-called cosmological constant (CC) Λ, introduced at first by Einstein in his gravitational field equations as a Lagrange multiplier ensuring the constancy of the 4-volume of the universe [13, 14]. However, CC explanation of dark energy encounters some fundamental obstacles in physics, in particular, the *fine-tuning* problem and the *coincidence* problem [15]. From the particle physics perspective, CC should be interpreted as the density of vacuum energy. The physical CC should contains quantum corrections from the zero-point energies of matter fields, which is close to Planck density M_P^4 ($M_P = 1/\sqrt{8\pi G}$ is the reduced Planck mass) in magnitude. What the fine-tuning problem has to face is the great discrepancy between the observed value of CC and its quantum correction, the former is only 10^{-123} times in magnitude as the latter [16, 17, 18, 19, 20, 21]. Even if this fine-tuning problem could be evaded, the coincidence problem as to why both energy densities of the observed dark energy and the dark matter are of the same order at present epoch remains, due to the fact that CC is time independent and non-dynamical.

The resolution of fine-tuning problem has probably to wait for the advent of a satisfied quantum theory of gravity in 4-dimensional spacetime. As a temporary expedient in cosmology community, the physical CC is assumed to vanish exactly[1] and it is conjectured that the dark energy which drives the late time cosmic accelerated expansion is dynamical. The observations actually say little about the evolution of the equation of state (EoS) of dark energy. Researchers have proposed lots of alternative models (see Reviews [23, 15, 24] and references therein) to explain the late time cosmic acceleration and alleviate the corresponding coincidence problem. The well-known Chaplygin gas model [25] and its generalizations, e.g., the models proposed in Refs. [26, 27, 28] are among these approaches.

---

[1] This assumption does actually prohibit the interpretation of CC as the energy density of vacuum fields. CC does not fluctuate, but the vacuum fields fluctuate and their fluctuations couple universally to gravitation [22]. We thank B. Deiss for pointing it to us.

The Chaplygin gas and its generalizations are of the so-called barotropic fluids whose pressure depends only upon the energy density [15]. What distinguishes them from other dynamic dark energy models is that, at background level, they get rid of the coincidence problem by unifying the dark energy and dark matter into a single dark substance. The gas behaves as a pressureless matter for very large redshift, however, it becomes a CC as the redshift is small. Due to the non-vanishing squared speed of sound, unfortunately, the Chaplygin gas and its generalizations would have fatal flaws for explaining tiny but stable CMB anisotropies [29, 30]. It follows from Sandvik's analysis in Ref. [30] that the unique reliable Chaplygin gas like model is the perfect fluid of constant pressure. That the pressure of a fluid is kept invariant makes the squared sound speed vanish identically, resulting in the disappearance of the embarrassed oscillation-instability issue in density perturbations. The density contrasts of constant-pressure barotropic fluid decay monotonously during evolution, implying that the fluid does not unify the dark energy and cold dark matter into a single dark substance. It is merely a mixture of CC and a non-relativistic matter, and is equivalent to the standard ΛCDM model to all orders in perturbation theory. Consequently, the coincidence problem remains.

The fluid models provide at most some effective descriptions of the dynamic dark energy. In view of the superstring/M-theory, a developing but promising candidate of quantum gravity, the realization of dark energy with some scalar fields is a more fundamental description than the perfect fluid models. The scalar fields are ubiquitous in superstring/M-theory, they arise naturally as dilaton or the compactification moduli. A scalar field with a canonical kinetic energy is dubbed *quintessence* [31], which is among the most studied candidates for dynamic dark energy. The quintessence typically involves a single scalar field with a particular self-interaction potential, allowing the vacuum energy to become dominant only recently. The quintessence potential has generally to be fine-tuned *ad hoc* to solve the coincidence problem. At background level, a quintessence field is generally equivalent to a perfect fluid whose EoS evolves from *plus one* at early times to approximately *minus one* at present epoch, with some unphysical ingredients introduced into the description of the evolution of Universe.

In this paper, we intend to study the cosmology of the quintessence counterpart of a perfect fluid of constant pressure. Different from Chaplygin gas like models, a quintessence field can not be viewed as a barotropic fluid in general [32]. Whether the quintessence field of constant pressure is equivalent to ΛCDM model or provides a unified description to dark energy and dark matter are open issues. Our main motivation in this paper is to establish a dynamic dark energy model with a quintessence scalar field whose EoS evolves from zero at early times to nearly minus one at present and potential is not required to be fine-tuned severely, so that the model is brought back to life in search of solutions of the coincidence problem. We also try to examine the possibility to build a unified description of dark energy and dark matter in such a scenario. It is shown that the constant pressure requirement enables the quintessence EoS interpolates between two plateaus, one corresponds to cold dark matter in the large redshift epoch, another to CC as the redshift comes near zero. The quintessence potential comes completely from the constant pressure assumption. It is factorized into the product of the squared Hubble rate and a dimensionless quantity, dubbed the *reduced* quintessence potential, which has also two plateaus so that the fine-tuning of the potential is unnecessary. Our investigation in the linear perturbation theory shows that the squared sound speeds of quintessence perturbations oscillate around *unity*, and the corresponding density contrasts decay monotonously during evolution. We conclude that the quintessence field of constant pressure could only play the role of a dynamic dark energy, which is neither a unified dark matter nor a mixture of CC and some non-relativistic matter.

Throughout the paper we adopt the Planck units $c = \hbar = M_P = 1$.

2 Model

We begin with the assumption that at the background level our universe is described by a flat Robertson-Walker metric

$$ds^2 = -dt^2 + a^2(t)d\vec{x}^2 \tag{1}$$

in which fills a mixture of a quintessence scalar $\phi(t)$ and a perfect fluid. The quintessence field is described by Lagrangian

$$\mathscr{L} = -\frac{1}{2}\partial_\mu\phi\partial^\mu\phi - V(\phi) \tag{2}$$

but the extra fluid by its pressure P_f and energy density ρ_f. There are several well-known candidates for such a cosmic fluid, *e.g.*, radiation, baryonic dust, the cold dark matter (CDM), the cosmological constant Λ, or mixture of them. As usual, we use ω_f to denote the EoS of this extra fluid, defined by $\omega_f = P_f/\rho_f$. The quintessence EoS is similarly defined as $\omega_\phi = P_\phi/\rho_\phi$, where $P_\phi = \frac{1}{2}\dot{\phi}^2 - V(\phi)$ and $\rho_\phi = \frac{1}{2}\dot{\phi}^2 + V(\phi)$ are respectively the pressure and energy density when the quintessence field is also viewed as a perfect fluid. $\omega_\phi \approx 1$ if kinetic term dominates while $\omega_\phi \approx -1$ if potential term dominates. If both kinetic and potential terms are almost equivalently important, $\omega_\phi \approx 0$. It is also remarkably for the EoS of a quintessence field not to cross over the cosmological constant boundary $\omega_{CC} = -1$ [33, 13, 24].

In the flat Robertson-Walker background, Einstein's gravitational field equations become the so-called Friedmann equations:

$$3H^2 = \frac{1}{2}\dot{\phi}^2 + V(\phi) + \rho_f \tag{3}$$

$$\dot{H} = -\frac{1}{2}\dot{\phi}^2 - \frac{1}{2}(1+\omega_f)\rho_f \tag{4}$$

where $H = \dot{a}/a$ is the Hubble's expansion rate of the universe for which we denote its present value as H_0. The extra fluid is assumed not to interact with the quintessence scalar field, $\dot{\rho}_f + 3H(1+\omega_f)\rho_f = 0$. Consequently, the consistency between two Friedmann equations requires that the evolution of the scalar field is subject to the following Klein-Gordon equation,

$$\ddot{\phi} + 3H\dot{\phi} + V_{,\phi} = 0 \tag{5}$$

To be convenient, we will use the so-called *efolding* number $N = \ln a$ as the time variable from now on, $-\infty < N < +\infty$, and let $N = 0$ represent the present epoch. A prime will represent a derivative with respect to N, unless otherwise specified. In terms of N, Eqs.(3), (4) and (5) are recast as:

$$3H^2 = V(\phi) + \frac{1}{2}H^2\phi'^2 + \rho_f \tag{6}$$

$$(H^2)' = -H^2\phi'^2 - (1+\omega_f)\,\rho_f \tag{7}$$

and

$$H^2\phi'' + \left[V(\phi) + \frac{1}{2}(1-\omega_f)\rho_f\right]\phi' + V_{,\phi}(\phi) = 0 \tag{8}$$

respectively.

We want to study a quintessence model in this paper whose EoS interpolates between the pressureless matter dominant epoch at large red shifts ($z \gtrsim 1$)[2] and the dark energy dominant epoch at $z \approx 0$. The merits of such a model are as follows. Firstly, there are few unphysical ingredients of EoS different from $\omega = 1/3, 0$ and $\omega = -1$ involved in the alleviation of the coincidence problem by a quintessence field. Besides, the model provides the possibility to unify the dark energy and dark matter into a single dark substance, the so-called *unified dark matter* or *quartessence* [34]. To define the model, we assume that the quintessence potential $V(\phi)$ takes a factorized form,

$$V(\phi) = 3H^2 U(\phi) \tag{9}$$

where the dimensionless quantity $U(\phi)$ is referred to as the reduced quintessence potential. With this assumption, we can translate Klein-Gordon equation (8) into,

$$H^2 \left[\frac{1}{2}\frac{d}{dN} + 3U + \frac{3}{2}(1-\omega_f)\Omega_f\right]\left[\phi'^2 - 6(1 - U - \Omega_f)\right] = 0 \tag{10}$$

where $\Omega_f = \rho_f / 3H^2$ is the reduced energy density of the extra fluid. The first Friedmann equation (6) can equivalently be expressed as,

$$\frac{1}{2}\phi'^2 = 3(1 - U - \Omega_f) \tag{11}$$

In terms of Eq.(11), Klein-Gordon equation Eq.(10) turns out to be an identity. This is indeed the case. As we have explained, among the two Friedmann equations (6), (7) and the Klein-Gordon equation (8), only two of them are independent.

What looks fascinating here is that Eq.(11) is similar to the characteristic equation of an instanton in $(1+1)$ Euclidean space if the *potential* $U(\phi)$ has more than one vacua [35, 36]. We want to know if the cosmological observations could put some restrictions on the possible forms of $U(\phi)$. With $U(\phi)$, the energy density and pressure of the quintessence field are rewritten as,

$$P_\phi = H^2 \left(\frac{1}{2}\phi'^2 - 3U\right) \tag{12}$$

$$\rho_\phi = H^2 \left(\frac{1}{2}\phi'^2 + 3U\right) \tag{13}$$

---

[2] The red-shift parameter z is defined as $z = e^{-N} - 1$. Therefore, $z \gtrsim 1$ corresponds to $N \lesssim -0.693$.

We can further simplify Eqs.(12) and (13) into:

$$\rho_\phi = 3H^2(1-\Omega_f) \tag{14}$$

$$P_\phi = 3H^2(1-2U-\Omega_f) \tag{15}$$

by exploiting Eq.(11). The cosmology of the model at the background level is determined completely by quintessence EoS ω_ϕ and the effective EoS parameter $\omega_{\text{eff}} = (P_\phi + P_f)/(\rho_\phi + \rho_f)$ of the mixture fluid. Obviously,

$$\omega_\phi = 1 - \frac{2U}{1-\Omega_f} \tag{16}$$

and

$$\omega_{\text{eff}} = 1 - 2U - (1-\omega_f)\Omega_f \tag{17}$$

Provided $\omega_{\text{eff}}|_{N\approx 0} < -1/3$, the late-time accelerated expansion of our universe occurs. Therefore, the magnitude of the reduced quintessence potential at the present epoch must obey the inequality: $\frac{2}{3} - \frac{1}{2}(1-\omega_f)\Omega_f^{(0)} < U|_{N\approx 0}$. For example, if the extra cosmic fluid is the non-relativistic matter, $\omega_f \approx 0$, $\Omega_f^{(0)} \approx 0.26$, we have $U|_{N\approx 0} \gtrsim 0.54$.

The interpolation of ω_ϕ between zero and minus one can be achieved if the reduced quintessence potential $U(\phi)$ has at least two plateaus, one is at $U \approx \frac{1}{2}[1-(1-\omega_f)\Omega_f]$ for the large red-shift epoch and another is at $U \approx 1 - \frac{1}{2}(1-\omega_f)\Omega_f$ as the red-shift parameter z approaches to zero. To this end, the quintessence field under consideration is defined to have a vanishing adiabatic sound speed $c^2_{\phi,\text{ad}} = \partial P_\phi/\partial\rho_\phi$. In other words, the pressure P_ϕ is supposed not to change during the evolution of the universe. Keeping P_ϕ invariant implies that,

$$\omega'_\phi = 3\omega_\phi(1+\omega_\phi) \tag{18}$$

Obviously, the expected plateaus in the $\omega_\phi \sim N$ curve are guaranteed by this condition, one is $\omega_\phi = 0$ and another is $\omega_\phi = -1$. Eq.(18) can be translated into the definition equation for the reduced quintessence potential,

$$U' + 3\big[(1-U)(1-2U) - (1-U+\omega_f U)\Omega_f\big] = 0 \tag{19}$$

The quintessence field is expected to play the role of dark energy at low red shifts to excite the late time cosmic acceleration. The magnitude of the reduced potential U must be larger than $\frac{1}{2}(1-\Omega_f)$ so that $P_\phi < 0$. For this scalar behaves as the pressureless non-relativistic matter at large red shifts, the Hubble rate H have to ascend sharply with respect to the increase of the red-shift parameter.

In the absence of the extra fluid, $\Omega_f = 0$, Eq.(19) can be solved analytically, with a closed form solution given below:

$$U(N) = \frac{1 - \alpha + (2\alpha - 1)e^{3N}}{2 - 2\alpha + (2\alpha - 1)e^{3N}} \tag{20}$$

where $\alpha := U|_{N=0}$ is an integration parameter which stands for the value of the reduced quintessence potential $U(\phi)$ at present. As pointed out previously, $\alpha > 2/3$. Because $\alpha \neq 0$, we see that: $1/2 \leq U(N) \leq 1$.

Substitution of Eq.(20) into Eqs.(11) and (7) leads to,

$$\phi(N) = \frac{2}{\sqrt{3}} \left[\tanh^{-1} \frac{1}{\sqrt{2(1-\alpha)}} - \tanh^{-1} \sqrt{\frac{2(1-\alpha) + (2\alpha - 1)e^{3N}}{2(1-\alpha)}} \right] \tag{21}$$

and

$$H^2 = H_0^2 \left[(2\alpha - 1) + 2(1 - \alpha)e^{-3N} \right] \tag{22}$$

To guarantee the positivity of H^2 during its evolution, $1/2 \leq \alpha \leq 1$[3]. Therefore, the parameter α takes its value in the region $2/3 < \alpha \leq 1$. Combination of Eqs.(21) and (22) with Eq.(9) enables us to express the quintessence potential as an explicit function of the scalar field itself. The result is,

$$V(\phi) = \frac{3H_0^2}{4} \left[3(2\alpha - 1) + (3 - 2\alpha)\cosh(\sqrt{3}\phi) - 2\sqrt{2(1-\alpha)}\sinh(\sqrt{3}\phi) \right] \tag{23}$$

There exist potentially three different interpretations of the quintessence field under consideration. The first possibility is simply to interpret it as a dynamic dark energy. Alternatively it could be interpreted as a unified dark matter (quartessence). With these two interpretations, the quintessence field is characterized by its energy density and pressure given below,

$$\rho_\phi = 3H_0^2 \left[(2\alpha - 1) + 2(1 - \alpha)e^{-3N} \right] \tag{24}$$

$$P_\phi = -3H_0^2(2\alpha - 1) \tag{25}$$

Though the pressure P_ϕ is a negative constant, the energy density ρ_ϕ interpolates between that of a non-relativistic matter, $\rho_\phi \approx 6H_0^2(1 - \alpha)e^{-3N}$, for

---

[3] When $\alpha = 1$, the quintessence scalar field under consideration degenerates to the cosmological constant $\Lambda = 3H_0^2$.

$N \ll 1$ and that of a cosmological constant, $\rho_\phi \approx 3H_0^2(2\alpha - 1)$, for $N \gg 0$. The evolution of quintessence EoS is found to be,

$$\omega_\phi = -\frac{(2\alpha - 1)e^{3N}}{2(1-\alpha) + (2\alpha - 1)e^{3N}} \tag{26}$$

In Figure 1, we plot the evolution of ω_ϕ by choosing $\alpha = 0.87$. As expected, ω_ϕ interpolates between minus one for $N \gtrsim 1$ and zero for $N \lesssim -2$.



Figure 1: Evolution of EoS ω_ϕ given in Eqs.(26) for parameters $\alpha = 0.87$. As expected, two plateaus exist. One plateau corresponds to $\omega_\phi \approx -1$ for $N \gtrsim 1$, another plateau corresponds to $\omega_\phi \approx 0$ for $N \lesssim -2$. ω_ϕ crosses over the acceleration boundary $-1/3$ at $N \approx -0.37$.

The third possibility is to interpret the quintessence field under consideration as a mixture of CC and some non-relativistic matter, $\rho_\phi = \rho_{CC} + \rho_M$. In this scenario, the dark energy is understood as a cosmological constant, $\rho_{CC} = -P_{CC} = 3H_0^2(2\alpha - 1)$, but the non-relativistic matter is characterized by $\rho_M = 6H_0^2(1-\alpha)e^{-3N}$ and $P_M = 0$. The EoS parameters of two components are $\omega_{CC} = -1$ and $\omega_M = 0$, respectively. Because of $\omega_{CC} = -1$, there is no interaction between the components. The quintessence EoS ω_ϕ, as given in Eq.(26), should be understood as the effective EoS of such a mixture, $\omega_\phi = \omega_{CC}\Omega_{CC} + \omega_M\Omega_M = -\Omega_{CC}$, where the dimensionless energy densities of the components are as follows,

$$\Omega_{CC}(N) = \frac{(2\alpha - 1)}{(2\alpha - 1) + 2(1-\alpha)e^{-3N}} \tag{27}$$

$$\Omega_M(N) = \frac{2(1-\alpha)e^{-3N}}{(2\alpha - 1) + 2(1-\alpha)e^{-3N}} \tag{28}$$

Notice that $\Omega_M(0) = 2(1-\alpha)$. Relying on the observational constraint $\Omega_M(0) \approx 0.26$, the best-fit value of the parameter α might be $\alpha \approx 0.87$. With respect to such an interpretation, the quintessence model under consideration is merely a reformulation of the standard ΛCDM model.

All of three interpretations of the quintessence field of a constant pressure appear to be plausible at the background level. They can, however, be distinguished from each other if we study the evolution of the matter perturbations. This will be addressed in the next section.



Figure 2: Parameter space of α and $\Omega_f^{(0)}$. The solid black lines stand for the critical cases, from bottom to the top $\alpha = (1-\Omega_f^{(0)})/2$, $\alpha = 1-\Omega_f^{(0)}$ and $\alpha = 1-\Omega_f^{(0)}/2$ respectively. The Dashed line corresponds to the acceleration boundary $6\alpha - 4 + 3\Omega_f^{(0)} = 0$. All of the parameters $(\Omega_f^{(0)}, \alpha)$ in the region above this dashed line allow the existence of late time cosmic acceleration.

When the contribution of the extra cosmic fluid is taken into account,

$$\Omega_f = \frac{H_0^2}{H^2}\Omega_f^{(0)} e^{-3N(1+\omega_f)} \tag{29}$$

Eqs. (7) and (11) become,

$$(H^2)' + 6H^2(1-U) - 3H_0^2\Omega_f^{(0)}(1-\omega_f)e^{-3N(1+\omega_f)} = 0 \tag{30}$$

$$\phi'^2 = 6(1-U) - 6\Omega_f^{(0)}\frac{H_0^2}{H^2}e^{-3N(1+\omega_f)} \tag{31}$$

respectively, and the condition (19) for preserving P_ϕ to be a constant during the evolution of the Universe is recast as:

$$U' + 3\left[(1-U)(1-2U) - \Omega_f^{(0)}\frac{H_0^2}{H^2}(1-U+\omega_f U)e^{-3N(1+\omega_f)}\right] = 0 \tag{32}$$

The closed-form solutions to Eqs.(30), (31) and (32) are not available in general. However, if the extra fluid is of extremely non-relativistic, $\omega_f = 0$, these equations can be solved exactly. The corresponding solution is as follows:

$$\phi(N) = \frac{2}{\sqrt{3}}\sqrt{\frac{2(1-\alpha-\Omega_f^{(0)})}{2-2\alpha-\Omega_f^{(0)}}}\left[\tanh^{-1}\frac{1}{\sqrt{2-2\alpha-\Omega_f^{(0)}}} - \tanh^{-1}\sqrt{\frac{2-2\alpha-\Omega_f^{(0)}+(2\alpha-1+\Omega_f^{(0)})e^{3N}}{2-2\alpha-\Omega_f^{(0)}}}\right] \tag{33}$$

$$H^2 = H_0^2\left[(2\alpha-1+\Omega_f^{(0)}) + (2-2\alpha-\Omega_f^{(0)})e^{-3N}\right] \tag{34}$$

$$U = \frac{1-\alpha-\Omega_f^{(0)}+(2\alpha-1+\Omega_f^{(0)})e^{3N}}{2-2\alpha-\Omega_f^{(0)}+(2\alpha-1+\Omega_f^{(0)})e^{3N}} \tag{35}$$

where as before, $\alpha = U|_{N=0}$. Substitution of Eqs.(35) and (29) into (16) and (17) leads to the following expressions of the quintessence EoS,

$$\omega_\phi = -\frac{(2\alpha-1+\Omega_f^{(0)})e^{3N}}{2(1-\alpha-\Omega_f^{(0)})+(2\alpha-1+\Omega_f^{(0)})e^{3N}} \tag{36}$$

and the effective EoS of the mixture,

$$\omega_{\text{eff}} = -\frac{(2\alpha-1+\Omega_f^{(0)})e^{3N}}{2-2\alpha-\Omega_f^{(0)}+(2\alpha-1+\Omega_f^{(0)})e^{3N}} \tag{37}$$

Manifestly, $\omega_\phi \lesssim \omega_{\text{eff}}$. Eqs.(34), (36) and (37) implies that the behavior of the mixed quintessence field and the extra fluid is determined by two parameters, i.e., the initial value of the reduced quintessence potential α and the initial dimensionless density $\Omega_f^{(0)}$ of the extra fluid. The late time accelerated expansion occurs if these two parameters are subject to the inequality $6\alpha - 4 + 3\Omega_f^{(0)} > 0$. In principle, α and $\Omega_f^{(0)}$ are two free parameters, however, there exist three degenerate but important cases where they are dependent upon one another. The first case corresponds to $\alpha = (1 - \Omega_f^{(0)})/2$, for which $\omega_\phi = \omega_{\text{eff}} = 0$, both of the quintessence field and the extra fluid act as non-relativistic matters. In the second case, $\alpha = 1 - \Omega_f^{(0)}/2$, $\omega_{\text{eff}} = -1$, the mixture of the quintessence field and the extra fluid behaves as a CC. The third degenerate case is characterized by equations $\alpha = 1 - \Omega_f^{(0)}$ and $\omega_\phi = -1$, in which the quintessence field itself is nothing but a CC. It is conspicuous that the standard ΛCDM is recovered in the third case if we take the values of the two parameters as $\alpha \approx 0.74$ and $\Omega_f^{(0)} \approx 0.26$.



Figure 3: Evolution of EoS ω_ϕ and ω_{eff} versus $N = \ln a$ for parameters $\alpha = 0.76$, $\omega_f = 0$ and $\Omega_f^{(0)} \approx 0.22$. The black curve stands for ω_{eff} while the gray curve describes ω_ϕ. Both curves interpolate between two plateaus of zero and of minus one.

If the extra fluid is assumed to describe the cold dark matter, e.g., $\Omega_f^{(0)} \approx 0.22$, $\alpha \approx 0.76$, ω_ϕ has remarkable differences from ω_{eff}. In particular, it is earlier for ω_ϕ to cross the acceleration boundary of $\omega = -1/3$. If the extra fluid is used to mimic all of the non-relativistic cosmic matter, i.e., the cold dark matter plus the baryonic dust, $\Omega_f^{(0)} \approx 0.26$ and $\alpha \approx 0.74$, ω_ϕ is almost the same as $\omega_{\rm eff}$. In all cases, both ω_ϕ and $\omega_{\rm eff}$ interpolate between two plateaus of zero and of minus one, so that the notorious coincidence problem with which the prototypic ΛCDM model is embarrassed is greatly alleviated in the quintessence model under consideration. The existence of plateaus in the $\omega \sim N$ curve implies that the severe fine-tuning of the parameters is unnecessary.

3 perturbations

Dark energy influences not only the expansion rate, it influences also the growth rate of matter perturbations. To clarify what the role does the quintessence field of constant pressure play during the evolution of the universe, in this section we study the matter perturbations at the linear perturbation level. We start with the perturbed Robertson-Walker metric,

$$ds^2 = -(1+2\Psi)H^{-2}dN^2 + e^{2N}(1+2\Phi)\delta_{ij}dx^i dx^j \tag{38}$$

in Newtonian gauge and for convenience let us work out from beginning a single Fourier mode k so that the perturbed quantities $\Phi = \Phi(N)e^{i\vec{k}\cdot\vec{x}}$, $\Psi = \Psi(N)e^{i\vec{k}\cdot\vec{x}}$ and so on. The perturbed energy-momentum tensor of the extra fluid is described by the perturbed energy density $\delta\rho_f = -T^0_{0(f)}$, the perturbed pressure $\delta P_f = \frac{1}{3}\delta T^i_{i(f)}$ and the velocity divergence θ_f defined by $ik_j\delta T^j_{0(f)} = -(1+\omega_f)\rho_f\theta_f$. For the quintessence field, these perturbed quantities read:

$$\delta\rho_\phi = H^2(\phi'\varphi' - \Psi\phi'^2) + V_{,\phi}\varphi \tag{39}$$

$$\delta P_\phi = H^2(\phi'\varphi' - \Psi\phi'^2) - V_{,\phi}\varphi \tag{40}$$

$$\theta_\phi = \hat{\lambda}^{-2}\frac{\varphi}{\phi'} \tag{41}$$

where $\hat{\lambda} = He^N/k$, and $\varphi := \delta\phi$ denotes the field fluctuation. The perturbation equations for a perfect fluid with density contrast δ_X and velocity divergence θ_X are:

$$\delta'_X = 3(\omega_X - c_X^2)\delta_X - (1+\omega_X)(\theta_X + 3\Phi') \tag{42}$$

$$\theta'_X = -\frac{\theta_X}{2}\left[1 - 6\omega_X - 3\omega_{\rm eff} + \frac{2\omega'_X}{1+\omega_X}\right] + \hat{\lambda}^{-2}\left[\frac{c_X^2\delta_X}{1+\omega_X} + \Psi\right] \tag{43}$$

where X is either ϕ or f, and $c_X^2 := \delta P_X/\delta\rho_X$ the squared sound speed of fluid X. The evolution of Bardeen potentials Φ and Ψ is subject to Einstein's gravitational equations. The result is,

$$\Phi = 3\hat{\lambda}^2\left[\frac{1}{2}\Omega_f\delta_f + \frac{1}{2}(1-\Omega_f)\delta_\phi + \Psi - \Phi'\right] \tag{44}$$

$$\Phi' = \Psi - \frac{3}{2}\hat{\lambda}^2\Big[(1+\omega_f)\Omega_f\theta_f + (1+\omega_\phi)(1-\Omega_f)\theta_\phi\Big] \tag{45}$$

$$\Psi = -\Phi \tag{46}$$

We assume that the extra fluid is of barotropic so that $\omega_f = \omega_f(\rho_f)$ and $c_f^2 = dP_f/d\rho_f$. The quintessence field, however, can not be regarded as a barotropic fluid in general. When X in Eqs.(42) and (43) stands for the quintessence field ϕ of constant pressure, ω_ϕ is given in Eq.(16), but the squared sound speed c_ϕ^2 does not coincide with its adiabatic counterpart $c_{\phi,\mathrm{ad}}^2$,

$$c_\phi^2 = \frac{H^2(\phi'\varphi' - \Psi\phi'^2) - V_{,\phi}\varphi}{H^2(\phi'\varphi' - \Psi\phi'^2) + V_{,\phi}\varphi} \tag{47}$$

In other words, c_ϕ^2 depends on the detailed behavior of both perturbation and background quantities. There is no reason for c_ϕ^2 vanishes identically. In fact, if we put ourselves in the quintessence rest frames, we have $\varphi = 0$ and hence $c_\phi^2 = 1$. Without loss of generality, we assume $c_\phi^2 \neq 0$ in the following. Eq.(43) for θ_ϕ turns out to become an identity, but Eq.(42) for δ_ϕ reduces to:

$$\varphi'' + \left(3 + \frac{H'}{H}\right)\varphi' + \left(\hat{\lambda}^{-2} + \frac{V_{,\phi\phi}}{H^2}\right)\varphi + 4\phi'\Phi' - 2\frac{V_{,\phi}}{H^2}\Phi = 0 \tag{48}$$

where,

$$V_{,\phi} = -\frac{3}{2}H^2\phi' \tag{49}$$

$$V_{,\phi\phi} = \frac{9}{4}H^2(3 - 2U - \Omega_f) \tag{50}$$

With respect to the observational constraints, the galaxy clustering on scales down to $10h^{-1}$Mpc requires the inequality $k \gtrsim 300H_0$ for the wave number of the perturbation modes. These modes are deep within the comoving horizon. Consequently, Eqs.(42), (43) for the extra fluid perturbations and Eq.(48) for the quintessence fluctuations should be solved in the sub-horizon limit, for which $\hat{\lambda} \ll 1$. Let us consider temporarily the case of $\Omega_f = 0$. Notice that $\Phi \sim \mathscr{O}(\hat{\lambda}^2)$ and $U \sim \mathscr{O}(1)$. The terms related to Φ, Φ' and $V_{,\phi\phi}/H^2$ in Eq.(48) are negligible with respect to the term involved in $\hat{\lambda}^{-2}$. Because $k \gg H_0$ on sub-horizon scales, the average of φ over many oscillations gives zero, $\langle\varphi\rangle \approx 0$. Consequently, the velocity divergence θ_ϕ of the quintessence mode oscillates around zero and its sound speed oscillates around $\langle c_\phi^2\rangle \approx 1$. The vanishing of the average velocity divergence $\langle\theta_\phi\rangle$ further implies the decoupling of Eq.(42) from Eq.(43) for $X = \phi$. The former turns out to become a first-order differential equation of δ_ϕ for $k \gg He^N$:

$$\delta'_\phi + 3(1 - \omega_\phi)\delta_\phi = 0 \tag{51}$$

The solution of Eq.(51) is stable and scale independent,

$$\delta_\phi(N) = \frac{\delta_\phi(0)e^{-3N}}{2(1-\alpha) + (2\alpha - 1)e^{3N}} \tag{52}$$

Unfortunately, such a density contrast decays monotonically, its contribution to the galaxy clustering should be negligible. It is the same as the sub-horizon density contrasts of the barotropic fluid of constant pressure [30]. Therefore, the quintessence field of constant pressure does not unify the dark energy and cold dark matter into a single dark substance. Notice that the sound speed of the mixture of CC and some non-relativistic matter vanishes identically [15]. The non-vanishing average $\langle c_\phi^2\rangle \approx 1$ which obviously mismatches this fact expels definitely the possibility to interpret the quintessence field of constant pressure as a mixture of CC and non-relativistic matters. The quintessence field of constant pressure describes only a dynamic dark energy. It clusters, but its clustering decays monotonously during the evolution and is therefore not directly responsible to the structure formation on large scales. What is in charge of the structure formation must have other candidates in the considered quintessence scenario.

The cold dark matter which is believed to take charge of the structure formation can effectively be described as an extremely non-relativistic perfect fluid. We should take $\Omega_f \neq 0$ and $\omega_f = 0$ in our model building. In such a scenario, the extra fluid is a mixture of cold dark matter and baryons. Although the solution to Eq.(48) gives still a rapid (damped) oscillating field fluctuation φ around $\langle\varphi\rangle = 0$ on deep sub-horizon scales so that the quintessence density contrasts decay rapidly, the density contrasts of the



Figure 4: Evolution of field fluctuation φ. It is obtained from the numerical solution of Eq.(48) for $k = 300H_0$. The other parameters are taken to be $\alpha = 0.87$ and $\Omega_f^{(0)} = 0$. The field fluctuation undergoes the damped oscillations around zero with a very high frequency.

non-relativistic fluid grow stably. It follows from Eqs.(42), (43), (44), (45) and (46) that, for $k \gg He^N$,

$$\delta'_\phi + 3(1-\omega_\phi)\delta_\phi = 0 \tag{53}$$

$$\delta''_f + \frac{1}{2}(1-3\omega_{\text{eff}})\delta'_f - \frac{3}{2}\Omega_f\delta_f = \frac{3}{2}(1-\Omega_f)\delta_\phi \tag{54}$$

where the fact $\theta_\phi \approx 0$ on deep sub-horizon scales has been made use of. Eq.(53), as expected, describes a decaying field density contrast δ_ϕ. Eq.(54), on the other hand, allows a growing density contrast δ_f for matter perturbations. It follows from Eq.(54) that the quintessence density contrast plays the role of an external source to the evolution of the density contrast of the non-relativistic fluid. If this external source is ignored, Eq.(54) can approximately be solved by a closed-form solution,

$$\begin{aligned} \delta_f(N) &= c_1 e^{3\gamma_- N}\ {}_2F_1(\gamma_-, \gamma_+ + 1/2, 1+\gamma_+ + \gamma_-; \xi) \\ &\quad + c_2 e^{-3\gamma_+ N}\ {}_2F_1(-\gamma_- + 1/2, -\gamma_+, 1-\gamma_+ - \gamma_-; \xi) \end{aligned} \tag{55}$$

where c_1 and c_2 are two integration constants,

$$\gamma_\pm = \frac{1}{12}\left[\sqrt{\frac{2\alpha - 23\Omega_f^{(0)} - 2}{2\alpha + \Omega_f^{(0)} - 2}} \pm 1\right], \qquad \xi = \frac{2\alpha + \Omega_f^{(0)} - 1}{2\alpha + \Omega_f^{(0)} - 1} e^{3N} \tag{56}$$

and ${}_2F_1(a,b,c;\xi)$ stands for the hypergeometric function. If $\alpha = 1 - \Omega_f^{(0)}$, for which the quintessence field degenerates to a CC (thereby $\delta_\phi = 0$), the solution given in Eq.(55) becomes of exact,

$$\delta_m(N) = c_1 e^N\ {}_2F_1\left(\frac{1}{3}, 1, \frac{11}{6}; \frac{\alpha e^{3N}}{\alpha - 1}\right) + c_2 e^{-3N/2}\ {}_2F_1\left(\frac{1}{6}, -\frac{1}{2}, \frac{1}{6}; \frac{\alpha e^{3N}}{\alpha - 1}\right) \tag{57}$$

The evolution of density contrast of non-relativistic matters in ΛCDM model is described by solution (57) for $\alpha \approx 0.74$. If the external source term is taken into account, a closed-form solution to Eq.(54) seems to be impossible. Nevertheless, the equation can be solved numerically if some appropriate initial conditions are assigned. The numerical solution of δ_f to Eq.(54) for initial conditions $\delta_f|_{N=-2} \approx 10^{-5}$ and $\delta_f'|_{N=-2} \approx 10^{-5}$ is plotted in Figure 5, where we have taken the values of the relevant parameters as $\alpha = 0.74$ and $\Omega_f^{(0)} = 0.26$. In the same figure we also plot the matter density contrast in ΛCDM model for comparison. The evolution of matter density contrasts in the two model does almost coincide with one another for $N \gtrsim -3$. When $N \lesssim -3$, the δ_f decays monotonically in our model but the ΛCDM density contrast grows monotonically. This difference implies probably that the growth rates of the matter perturbations are different in two models.

The growth rate of matter perturbation is defined as the ratio of the derivative of the matter density contrast with respect to enfolding time to matter density contrast itself, which reads $f = \delta_f'/\delta_f$ in our model. In Figure 6, we plot the evolution of the growth rate of matter perturbations in our model as well as its analogue in ΛCDM. Because the cosmological constant does not cluster, it has no influence on the evolution of matter perturbations, the growth rate of matter perturbations decays monotonically in the standard ΛCDM model. If the late time acceleration is motivated by some dynamic dark energy, e.g., the quintessence scalar field of constant pressure, the clustering of dark energy perturbations that is generally inevitable would affect the growth of matter perturbations. In the proposed scenario, although the quintessence density contrast does not take charge of the large scale structures, it acts as an external driving force in the evolution equation (54) of matter density contrast. Such an influence results in a crest in $f(N)$ curve at $N \approx -3$, as plotted in Figure 6. When $N \lesssim -3$, the quintessence density contrast δ_ϕ has a sufficient magnitude to accelerate the growth of matter perturbations. When $N \gtrsim -3$, the magnitude of δ_ϕ drops effectively to zero and no longer provides robust driving force, so the growth rate of matter



Figure 5: Evolution of density contrast of matter perturbations on deep sub-horizon scales. The black curve describes the density contrast δ_f as the numerical solution of Eq.(54) for $\alpha = 0.74$ and $\Omega_f^{(0)} = 0.26$. The gray curve describes the density contrast in the standard ΛCDM model. The two curves coincide with each other with a very high accuracy for $N \gtrsim -3$.

perturbations decelerates.

4 Conclusion

In this paper we have studied the cosmology of a quintessence scalar field which can be viewed as a non-barotropic perfect fluid of constant pressure in linear perturbation theory. The assumption of constant pressure ensures the interpolation of the quintessence EoS between the plateau of zero at large red shifts and that of minus one at small red shifts. Not this characteristic alleviates greatly the coincidence problem only, it bring also few unphysical ingredients into the description of evolution of universe. The potential of the quintessence field is determined completely from the requirement of constant pressure, which is unnecessary to be severely fine-tuned in the alleviation of coincidence problem. The adiabatic sound speed of the quintessence field vanishes, however, its physical sound speed depends upon the details of perturbations and equals approximately to unity for modes within the deep comoving horizon. The quintessence field does not unify the cold dark matter



Figure 6: Evolution of growth rates of matter perturbations in the model under consideration and in ΛCDM model. The black curve describes the growth rate of the density contrast δ_f in our model as the numerical solution of Eq.(54) for $\alpha = 0.74$ and $\Omega_f^{(0)} = 0.26$. The gray curve is the growth rate of the matter density contrast in ΛCDM model. The two curves coincide for $N \gtrsim -3$ but differ clearly from one another for $N < -3$.

and dark energy into a single dark substance, it is also not a mixture of these two dark substances. What the role it really play during evolution is plausibly a dynamic dark energy that could cluster. This clustering is measured by a decaying density contrast. Though it is not responsible to the formation of large scale structure, the clustering of the quintessence perturbations impacts the developments of matter perturbations as an external driving force.

Acknowledgments

We would like to thank Y. F. Cai, M. Z. Li, J.X. Lu and M.L. Yan for valuable discussions. The work was supported in part by NSFC under No. 11235010.

References

[1] Adam G Riess, Alexei V Filippenko, Peter Challis, Alejandro Clocchiatti, Alan Diercks, M Peter, Ron L Gilliland, Craig J Hogan, Saurabh

Jha, Robert P Kirshner, and B Leibundgut. Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *Astron. J.*, 116:1009, 1998.

[2] S. Perlmutter, G. Aldering, G. Goldhaber, R. A. Knop, P. Nugent, P. G. Castro, S. Deustua, S. Fabbro, A. Goobar, D. E. Groom, I. M. Hook, A. G. Kim, M. Y. Kim, J. C. Lee, N. J. Nunes, R. Pain, C. R. Pennypacker, R. Quimby, C. Lidman, R. S. Ellis, M. Irwin, R. G. McMahon, P. Ruizapuente, N. Walton, B. Schaefer, B. J. Boyle, A. V. Filippenko, T. Matheson, A. S. Fruchter, N. Panagia, H. J. M. Newberg, W. J. Couch, and The Supernova Cosmology Project. Measurements of Ω and Λ from 42 High redshift Supernovae. *The Astrophysical Journal*, 517(2):565–586, June 1999.

[3] Max Tegmark, Michael Strauss, Michael Blanton, Kevork Abazajian, Scott Dodelson, Havard Sandvik, Xiaomin Wang, David Weinberg, Idit Zehavi, Neta Bahcall, Fiona Hoyle, David Schlegel, Roman Scoccimarro, Michael Vogeley, Andreas Berlind, Tamás Budavari, Andrew Connolly, Daniel Eisenstein, Douglas Finkbeiner, Joshua Frieman, James Gunn, Lam Hui, Bhuvnesh Jain, David Johnston, Stephen Kent, Huan Lin, Reiko Nakajima, Robert Nichol, Jeremiah Ostriker, Adrian Pope, Ryan Scranton, Uroš Seljak, Ravi Sheth, Albert Stebbins, Alexander Szalay, István Szapudi, Yongzhong Xu, James Annis, J. Brinkmann, Scott Burles, Francisco Castander, Istvan Csabai, Jon Loveday, Mamoru Doi, Masataka Fukugita, Bruce Gillespie, Greg Hennessy, David Hogg, Željko Ivezić, Gillian Knapp, Don Lamb, Brian Lee, Robert Lupton, Timothy McKay, Peter Kunszt, Jeffrey Munn, Liam Oonnell, John Peoples, Jeffrey Pier, Michael Richmond, Constance Rockosi, Donald Schneider, Christopher Stoughton, Douglas Tucker, Daniel Vanden Berk, Brian Yanny, and Donald York. Cosmological parameters from SDSS and WMAP. *Physical Review D*, 69(10):103501, May 2004.

[4] Max Tegmark, Michael R. Blanton, Michael A. Strauss, Fiona Hoyle, David Schlegel, Roman Scoccimarro, Michael S. Vogeley, David H. Weinberg, Idit Zehavi, Andreas Berlind, Tamas Budavari, Andrew Connolly, Daniel J. Eisenstein, Douglas Finkbeiner, Joshua A. Frieman, James E. Gunn, Andrew J. S. Hamilton, Lam Hui, Bhuvnesh Jain, David Johnston, Stephen Kent, Huan Lin, Reiko Nakajima, Robert C. Nichol, Jeremiah P. Ostriker, Adrian Pope, Ryan Scranton, Uroš Seljak,

Ravi K. Sheth, Albert Stebbins, Alexander S. Szalay, Istvan Szapudi, Licia Verde, Yongzhong Xu, James Annis, Neta A. Bahcall, J. Brinkmann, Scott Burles, Francisco J. Castander, Istvan Csabai, Jon Loveday, Mamoru Doi, Masataka Fukugita, J. Richard Gott III, Greg Hennessy, David W. Hogg, Željko Ivezić, Gillian R. Knapp, Don Q. Lamb, Brian C. Lee, Robert H. Lupton, Timothy A. McKay, Peter Kunszt, Jeffrey A. Munn, Liam Oonnell, John Peoples, Jeffrey R. Pier, Michael Richmond, Constance Rockosi, Donald P. Schneider, Christopher Stoughton, Douglas L. Tucker, Daniel E. Vanden Berk, Brian Yanny, and Donald G. York. The Three dimensional Power Spectrum of Galaxies from the Sloan Digital Sky Survey. *The Astrophysical Journal*, 606(2):702–740, May 2004.

[5] Max Tegmark, Daniel J. Eisenstein, Michael a. Strauss, David H. Weinberg, Michael R. Blanton, Joshua a. Frieman, Masataka Fukugita, James E. Gunn, Andrew J. S. Hamilton, Gillian R. Knapp, Robert C. Nichol, Jeremiah P. Ostriker, Nikhil Padmanabhan, Will J. Percival, David J. Schlegel, Donald P. Schneider, Roman Scoccimarro, Uroš Seljak, Hee-Jong Seo, Molly Swanson, Alexander S. Szalay, Michael S. Vogeley, Jaiyul Yoo, Idit Zehavi, Kevork Abazajian, Scott F. Anderson, James Annis, Neta a. Bahcall, Bruce Bassett, Andreas Berlind, Jon Brinkmann, Tamás Budavari, Francisco Castander, Andrew Connolly, Istvan Csabai, Mamoru Doi, Douglas P. Finkbeiner, Bruce Gillespie, Karl Glazebrook, Gregory S. Hennessy, David W. Hogg, Željko Ivezić, Bhuvnesh Jain, David Johnston, Stephen Kent, Donald Q. Lamb, Brian C. Lee, Huan Lin, Jon Loveday, Robert H. Lupton, Jeffrey a. Munn, Kaike Pan, Changbom Park, John Peoples, Jeffrey R. Pier, Adrian Pope, Michael Richmond, Constance Rockosi, Ryan Scranton, Ravi K. Sheth, Albert Stebbins, Christopher Stoughton, István Szapudi, Douglas L. Tucker, Daniel E. Vanden Berk, Brian Yanny, and Donald G. York. Cosmological constraints from the SDSS luminous red galaxies. *Physical Review D*, 74(12):123507, December 2006.

[6] Uroš Seljak, Alexey Makarov, Patrick McDonald, Scott Anderson, Neta Bahcall, J. Brinkmann, Scott Burles, Renyue Cen, Mamoru Doi, James Gunn, Željko Ivezić, Stephen Kent, Jon Loveday, Robert Lupton, Jeffrey Munn, Robert Nichol, Jeremiah Ostriker, David Schlegel, Donald Schneider, Max Tegmark, Daniel Berk, David Weinberg, and Donald

York. Cosmological parameter analysis including SDSS Lyα forest and galaxy bias: Constraints on the primordial spectrum of fluctuations, neutrino mass, and dark energy. *Physical Review D*, 71(10):103515, May 2005.

[7] Jennifer K. Adelman-McCarthy, J. K.cCarthy, Marcel A. Agueros, Sahar S. Allam, Kurt S. J. Anderson, Scott F. Anderson, James Annis, Neta A. Bahcall, Ivan K. Baldry, J. C. Barentine, Andreas Berlind, Mariangela Bernardi, Michael R. Blanton, William N. Boroski, Howard J. Brewington, Jarle Brinchmann, J. Brinkmann, Robert J. Brunner, Tamas Budavari, Larry N. Carey, Michael A. Carr, Francisco J. Castander, A. J. Connolly, Istvan Csabai, Paul C. Czarapata, Julianne J. Dalcanton, Mamoru Doi, Feng Dong, Daniel J. Eisenstein, Michael L. Evans, Xiaohui Fan, Douglas P. Finkbeiner, Scott D. Friedman, Joshua A. Frieman, Masataka Fukugita, Bruce Gillespie, Karl Glazebrook, Jim Gray, Eva K. Grebel, James E. Gunn, Vijay K. Gurbani, Ernst de Haas, Patrick B. Hall, Frederick H. Harris, Michael Harvanek, Suzanne L. Hawley, Jeffrey Hayes, John S. Hendry, Gregory S. Hennessy, Robert B. Hindsley, Christopher M. Hirata, Craig J. Hogan, David W. Hogg, Donald J. Holmgren, Jon A. Holtzman, Shinchi Ichikawa, Željko Ivezić, Sebastian Jester, David E. Johnston, Anders M. Jorgensen, Mario Jurić, Stephen M. Kent, S. J. Kleinman, G. R. Knapp, Alexei Yu. Kniazev, Richard G. Kron, Jurek Krzesinski, Nikolay Kuropatkin, Donald Q. Lamb, Hubert Lampeitl, Brian C. Lee, R. French Leger, Huan Lin, Daniel C. Long, Jon Loveday, Robert H. Lupton, Bruce Margon, David Martinezelgado, Rachel Mandelbaum, Takahiko Matsubara, Peregrine M. McGehee, Timothy A. McKay, Avery Meiksin, Jeffrey A. Munn, Reiko Nakajima, Thomas Nash, Eric H. Neilsen, Jr., Heidi Jo Newberg, Peter R. Newman, Robert C. Nichol, Tom Nicinski, Maria Nietoantisteban, Atsuko Nitta, William Oullane, Sadanori Okamura, Russell Owen, Nikhil Padmanabhan, George Pauls, John Peoples, Jr., Jeffrey R. Pier, Adrian C. Pope, Dimitri Pourbaix, Thomas R. Quinn, Gordon T. Richards, Michael W. Richmond, Constance M. Rockosi, David J. Schlegel, Donald P. Schneider, Joshua Schroeder, Ryan Scranton, Uroš Seljak, Erin Sheldon, Kazu Shimasaku, J. Allyn Smith, Vernesa Smolčić, Stephanie A. Snedden, Chris Stoughton, Michael A. Strauss, Mark SubbaRao, Alexander S. Szalay, Istvan Szapudi, Paula Szkody, Max Tegmark, Aniruddha R. Thakar, Douglas L. Tucker, Alan

Uomoto, Daniel E. Vanden Berk, Jan Vandenberg, Michael S. Vogeley, Wolfgang Voges, Nicole P. Vogt, Lucianne M. Walkowicz, David H. Weinberg, Andrew A. West, Simon D. M. White, Yongzhong Xu, Brian Yanny, D. R. Yocum, Donald G. York, Idit Zehavi, Stefano Zibetti, and Daniel B. Zucker. The Fourth Data Release of the Sloan Digital Sky Survey. *The Astrophysical Journal Supplement Series*, 162(1):38–48, January 2006.

[8] Kevork Abazajian, Jennifer K. Adelman-McCarthy, Marcel A. Agüeros, Sahar S. Allam, Kurt S. J. Anderson, Scott F. Anderson, James Annis, Neta A. Bahcall, Ivan K. Baldry, Steven Bastian, Andreas Berlind, Mariangela Bernardi, Michael R. Blanton, Jr. Bochanski, William N. Boroski, Howard J. Brewington, John W. Briggs, J. Brinkmann, Robert J. Brunner, Tamás Budavári, Larry N. Carey, Francisco J. Castander, A. J. Connolly, Kevin R. Covey, István Csabai, Julianne J. Dalcanton, Mamoru Doi, Feng Dong, Daniel J. Eisenstein, Michael L. Evans, Xiaohui Fan, Douglas P. Finkbeiner, Scott D. Friedman, Joshua A. Frieman, Masataka Fukugita, Bruce Gillespie, Karl Glazebrook, Jim Gray, Eva K. Grebel, James E. Gunn, Vijay K. Gurbani, Patrick B. Hall, Masaru Hamabe, Daniel Harbeck, Frederick H. Harris, Hugh C. Harris, Michael Harvanek, Suzanne L. Hawley, Jeffrey Hayes, Timothy M. Heckman, John S. Hendry, Gregory S. Hennessy, Robert B. Hindsley, Craig J. Hogan, David W. Hogg, Donald J. Holmgren, Jon A. Holtzman, Shinichi Ichikawa, Takashi Ichikawa, Željko Ivezić, Sebastian Jester, David E. Johnston, Anders M. Jorgensen, Mario Jurić, Stephen M. Kent, S. J. Kleinman, G. R. Knapp, Alexei Yu. Kniazev, Richard G. Kron, Jurek Krzesinski, Donald Q. Lamb, Hubert Lampeitl, Brian C. Lee, Huan Lin, Daniel C. Long, Jon Loveday, Robert H. Lupton, Ed Mannery, Bruce Margon, David Martínez-Delgado, Takahiko Matsubara, Peregrine M. McGehee, Timothy A. McKay, Avery Meiksin, Brice Ménard, Jeffrey A. Munn, Thomas Nash, Jr. Neilsen, Heidi Jo Newberg, Peter R. Newman, Robert C. Nichol, Tom Nicinski, Maria Nieto-Santisteban, Atsuko Nitta, Sadanori Okamura, William O'Mullane, Russell Owen, Nikhil Padmanabhan, George Pauls, John Peoples, Jeffrey R. Pier, Adrian C. Pope, Dimitri Pourbaix, Thomas R. Quinn, M. Jordan Raddick, Gordon T. Richards, Michael W. Richmond, Hans-Walter Rix, Constance M. Rockosi, David J. Schlegel, Donald P. Schneider, Joshua Schroeder, Ryan Scranton, Maki Sekiguchi, Erin Sheldon, Kazu Shimasaku, Nicole M.

Silvestri, J. Allyn Smith, Vernesa Smolčić, Stephanie A. Snedden, Albert Stebbins, Chris Stoughton, Michael A. Strauss, Mark SubbaRao, Alexander S. Szalay, István Szapudi, Paula Szkody, Gyula P. Szokoly, Max Tegmark, Luis Teodoro, Aniruddha R. Thakar, Christy Tremonti, Douglas L. Tucker, Alan Uomoto, Daniel E. Vanden Berk, Jan Vandenberg, Michael S. Vogeley, Wolfgang Voges, Nicole P. Vogt, Lucianne M. Walkowicz, Shu-i Wang, David H. Weinberg, Andrew A. West, Simon D. M. White, Brian C. Wilhite, Yongzhong Xu, Brian Yanny, Naoki Yasuda, Ching-Wa Yip, D. R. Yocum, Donald G. York, Idit Zehavi, Stefano Zibetti, and Daniel B. Zucker. The Third Data Release of the Sloan Digital Sky Survey. *The Astronomical Journal*, 129(3):1755–1759, March 2005.

[9] D. N. Spergel, R. Bean, O. Dore, M. R. Nolta, C. L. Bennett, J. Dunkley, G. Hinshaw, N. Jarosik, E. Komatsu, L. Page, H. V. Peiris, L. Verde, M. Halpern, R. S. Hill, A. Kogut, M. Limon, S. S. Meyer, N. Odegard, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright. Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology. *The Astrophysical Journal Supplement Series*, 170(2):377–408, June 2007.

[10] L. Page, G. Hinshaw, E. Komatsu, M. R. Nolta, D. N. Spergel, C. L. Bennett, C. Barnes, R. Bean, O. Dore, J. Dunkley, M. Halpern, R. S. Hill, N. Jarosik, A. Kogut, M. Limon, S. S. Meyer, N. Odegard, H. V. Peiris, G. S. Tucker, L. Verde, J. L. Weiland, E. Wollack, and E. L. Wright. Three year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Polarization Analysis. *The Astrophysical Journal Supplement Series*, 170(2):335–376, June 2007.

[11] G. Hinshaw, M. R. Nolta, C. L. Bennett, R. Bean, O. Dore, M. R. Greason, M. Halpern, R. S. Hill, N. Jarosik, A. Kogut, E. Komatsu, M. Limon, N. Odegard, S. S. Meyer, L. Page, H. V. Peiris, D. N. Spergel, G. S. Tucker, L. Verde, J. L. Weiland, E. Wollack, and E. L. Wright. Three year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Temperature Analysis. *The Astrophysical Journal Supplement Series*, 170(2):288–334, June 2007.

[12] N. Jarosik, C. Barnes, M. R. Greason, R. S. Hill, M. R. Nolta, N. Odegard, J. L. Weiland, R. Bean, C. L. Bennett, O. Dore, M. Halpern,

G. Hinshaw, A. Kogut, E. Komatsu, M. Limon, S. S. Meyer, L. Page, D. N. Spergel, G. S. Tucker, E. Wollack, and E. L. Wright. Three year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Beam Profiles, Data Processing, Radiometer Characterization, and Systematic Error Limits. *The Astrophysical Journal Supplement Series*, 170(2):263–287, June 2007.

[13] Martin Kunz and Domenico Sapone. Crossing the phantom divide. *Physical Review D*, 74:123503, 2006.

[14] Stefan Nobbenhuis. The cosmological constant problem, an inspiration for new physics. *Arxiv preprint gr-qc/0609011*, 2006.

[15] L Amendola and S. Tsujikawa. *Dark Energy, Theory and Observations.* CUP, 2010.

[16] Shamit Kachru, Michael Schulz, and Eva Silverstein. Self-tuning flat domain walls in 5D gravity and string theory. *Physical Review D*, 62:045021, 2000.

[17] S.-H. Henry Tye and Ira Wasserman. Brane World Solution to the Cosmological Constant Problem. *Physical Review Letters*, 86(9):1682–1685, February 2001.

[18] Jun'ichi Yokoyama. Cosmological Constant From Degenerate Vacua. *Physical Review Letters*, 88(15):8, March 2002.

[19] G. L. Kane, M. J. Perry, and A. N. Zytkow. A Possible Mechanism for Generating a Small Positive Cosmological Constant. *arXiv:hep-th/0311152*, 2003.

[20] Shinji Mukohyama and Lisa Randall. Dynamical Approach to the Cosmological Constant. *Physical Review Letters*, 92:4, 2004.

[21] A. Dolgov and F. Urban. Dynamical vacuum energy via adjustment mechanism. *Physical Review D*, 77:083503, 2008.

[22] Bruno M Deiss. Cosmic Dark Energy Emerging from Gravitationally Effective Vacuum Fluctuations. *arXiv:1209.5386*, 2012.

[23] Edmund J. Copeland, M. Sami, and Shinji Tsujikawa. Dynamics of dark energy. *International Journal of Modern Physics*, D15:1753–1936, 2006.

[24] Yi-Fu Cai, Emmanuel N. Saridakis, Mohammad R. Setare, and Jun-Qing Xia. Quintom cosmology: Theoretical implications and observations. *Physics Reports*, 493:1–60, 2010.

[25] Alexander Kamenshchik, Ugo Moschella, and Vincent Pasquier. An alternative to quintessence. *Physics Letters B*, 511(2-4):265–268, July 2001.

[26] M. Bento, O. Bertolami, and a. Sen. Generalized Chaplygin gas, accelerated expansion, and dark energy-matter unification. *Physical Review D*, 66:043507, 2002.

[27] Neven Bilić, GaryB Tupper, and RaoulD Viollier. Unification of dark matter and dark energy: the inhomogeneous Chaplygin gas. *Physics Letters B*, 535:17, 2002.

[28] Neven Bilić, Gary Tupper, and Raoul Viollier. Cosmological tachyon condensation. *Physical Review D*, 80:023515, 2009.

[29] L Amendola, F Finelli, C Burigana, and D Carturan. WMAP and the generalized Chaplygin gas. *Journal of Cosmology and Astroparticle Physics*, 2003:005–005, 2003.

[30] Havard Sandvik, Max Tegmark, Matias Zaldarriaga, and Ioav Waga. The end of unified dark matter? *Physical Review D*, 69:123524, 2004.

[31] Bharat Ratra and P. J Peebles. Cosmological Consequences of a Rolling Homogeneous Scalar Field. *Phys. Rev. D*, 37:3406, 1988.

[32] Adam J. Christopherson and Karim a. Malik. The non-adiabatic pressure in general scalar field systems. *Physics Letters B*, 675:159–163, 2009.

[33] Alexander Vikman. Can dark energy evolve to the phantom? *Physical Review D*, 71:023515, 2005.

[34] Martin Makler, Sergio Quinet de Oliveira, and Ioav Waga. Constraints on the generalized chaplygin gas from supernovae observations. *Physics Letters B*, 555:1, 2003.

[35] R. Rajaraman. *Solitons and instantons.* Amsterdam: North-Holland, 1982.

[36] T. Vachaspati. *Kinks and domain walls*. CUP, 2006.